Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

rather poor character of the plates of 1899 makes a definite decision impossible.

The star T *Tauri* itself has a very interesting nebulous wing which is either variable or must rotate so as to be concealed at times. This appendage is a small cone-shaped projection perhaps four seconds of arc in length in position angle 150 ± °. It was often seen by TEMPEL, and was fairly conspicuous when examined by BARNARD and BURNHAM in 1890, but entirely invisible when BARNARD looked for it in February, 1895. It can be clearly made out in the elongated image of T *Tauri* on both the Crossley plates of 1899. On the 1914 plates it is certainly not present; the disk of the variable is perfectly round, tho perhaps with just a trace of nebulosity all around it. It is evident that this interesting variable will repay further investigation.

H. D. CURTIS.

SEARCH FOR FAINT MEMBERS OF THE TAURUS CLUSTER.

Dr. HERTZSPRUNG suggested in 1914 that it would be of interest to compare the earlier plates of the region of T *Tauri* with recent plates in the endeavor to find very faint members of the well-known Taurus stream. This region is about five degrees from the center of the *Hyades,* and a proper motion of $1''.6$ would be expected in the interval of nearly fifteen years. A motion of this amount should show clearly on the Crossley plates.

The measures were made entirely differential, with respect to a common system of five stars of eighth to ninth magnitude; and twelve faint stars of from fourteenth to eighteenth magnitudes were selected at random in the area of best definition. None of the small relative motions found agree either in magnitude or direction with the motion of the components of the Taurus stream. Within the limited area examined the results must be regarded as purely negative.

H. D. CURTIS.

NOTE ON COMET *d* 1915 (MELLISH).

Comet *d* 1915 was first seen by Mr. JOHN E. MELLISH, of the Yerkes Observatory, on September 13, 1915.

The first observation received at the Students' Observatory